PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



(51) International Patent Classification 7:		11) International Publication Number:	WO 00/25766
A61K 31/00	A2		110 00/25/00
		43) International Publication Date:	11 May 2000 (11.05.00
 (21) International Application Number: PCT/GB (22) International Filing Date: 26 October 1999 ((30) Priority Data: 60/106,354 30 October 1998 (30.10.98) (71) Applicant (for all designated States except US): LIMITED [GB/GB]; 15 Stanhope Gate, London V (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): LENGEL, Date [US/US]; 1800 Concord Pike, P.O. Box 15437, William, Leroy 1800 Concord Pike, P.O. Box 15437, William, Leroy 1850—5437 (US). (74) Agent: BRYANT, Tracey; Global Intellectual Propertical Pr	ZENEC W1Y 6L wid, Jo ilmingto [US/US gton, I	(81) Designated States: AE, AL, AM, AT, BR, BY, CA, CH, CN, CR, CU, ES, FI, GB, GD, GE, GH, GM, HR KE, KG, KP, KR, KZ, LC, LK, LR MD, MG, MK, MN, MW, MX, NO, SD, SE, SG, SI, SK, SL, TJ, TM, US, UZ, VN, YU, ZA, ZW, ARIPLS, MW, SD, SL, SZ, TZ, UG, ZW, AZ, BY, KG, KZ, MD, RU, TJ, TN, BE, CH, CY, DE, DK, ES, FI, FR, MC, NL, PT, SE), OAPI patent (B, GA, GN, GW, ML, MR, NE, SN, SP, GM, GW, ML, MR, NE, SN, SP, GW, GW, ML, MR, NE, SN, SP, GW, GW, GW, GW, ML, MR, NE, SN, SP, GW, GW, GW, GW, GW, GW, GW, GW, GW, GW	CZ, DE, DK, DM, EB, HU, ID, IL, IN, IS, JF, LU, LV, MAD, NZ, PL, PT, RO, RU, TR, TT, TZ, UA, UCO patent (GH, GM, KE), Eurasian patent (AM), European patent (ATR, GB, GR, IE, IT, LUF, BJ, CF, CG, CI, CM, TD, TG).
(54) Title: TREATMENT OF GASTRIC ASTHMA			
Use of neurokinin 2 (NK2) receptor antagonists in the with gastroesophageal reflux, including pharmaceutical corcompositions.	e treatm	t or prevention of gastric asthma or adverse response containing the NK2 receptor antagonist and r	piratory events associate methods for making suc

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

of America

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	us	United States of Ameri
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

. -1-

TREATMENT OF GASTRIC ASTHMA

Background

10

15

20

25

30

Gastric asthma, described by Bruno et al. (Int. J. IMPH. 10/3: 195-204, 1997; Mays, E.E. JAMA 236: 2626, 1976), is a syndrome arising from the gastroesophageal acid induced changes in respiratory function. There is a progressive body of evidence supporting the role of gastroesophageal reflux as a triggering factor of bronchial hyperreactivity leading to this syndrome.

The mammalian neurokinins comprise a class of peptide neurotransmitters which are found in the peripheral and central nervous systems. The three principal neurokinins are Substance P (SP), Neurokinin A (NKA) and Neurokinin B (NKB). There are also N-terminally extended forms of at least NKA. At least three receptor types are known for the three principal neurokinins. Based upon their relative selectivities favoring the neurokinin agonists SP, NKA and NKB, the receptors are classifed as neurokinin 1 (NK1), neurokinin 2 (NK2) and neurokinin 3 (NK3) receptors, respectively. In the periphery, SP and NKA are localized in C-afferent sensory neurons, which neurons are characterized by non-myelinated nerve endings known as C-fibers, and are released by selective depolarization of these neurons, or selective stimulation of the C-fibers. C-Fibers are located in the airway epithelium, and the tachykinins are known to cause profound effects which clearly parallel many of the symptoms observed in asthmatics. The effects of release or introduction of tachykinins in mammalian airways include bronchoconstriction, increased microvascular permeability, vasodilation, increased mucus secretion and activation of mast cells. Thus, the tachykinins are implicated in the pathophysiology and airway hyperresponsiveness observed in asthmatics; and blockade of the action of released tachykinins may be useful in the treatment of asthma and related conditions.

Accordingly, NK2 antagonists have been implicated in the pathology of numerous diseases including asthma, airway oedema, bladder hypermotility, hypertension, pain, gastrointestinal-hypermotility, psychoses including schizophrenia and other psychatric indications, such as depression and anxiety.

The development of antagonists for the NK2 receptor has been purported to be useful for treatment of bronchial reactivity (Barnes, P.J. Lancet, i: 242-245, 1986). Although the etiological relationship between gastroesophageal reflux and asthma is not clear, the number

-2-

of reflux episodes can be correlated to those of bronchial hyperreactivity (Vincent et al., Eur. Respir. J. 10/10: 2255-2259, 1997). Current asthma therapies have not been shown to reverse bronchial hyperreactivity and in some cases are associated with adverse side effects (steroids), the progressive diminution of the therapeutic benefits (beta agonists) and they have not been shown to have any beneficial effects for gastroesophageal reflux induced respiratory changes. Moreover, chronic asthma treatments such as theophylline and beta agonists may promote or worsen gastroesophageal reflux (Bruno et al., Int. J. IMPH. 10/3: 195-204, 1997). Therefore, there is an unmet medical need for a treatment that interrupts the cyclic nature of gastroesophageal reflux and asthma.

There are no suggestions that NK2 antagonists could be beneficial in treating gastric asthma and in particular in treating gastric asthma associated with gastroesophageal reflux, which is manifested in adverse respiratory events such as chronic cough and asthma. The use of NK2 antagonists minimises the possibility of side-effects associated with alternative methods of treating such disease conditions and offers the physician a valuable alternative treatment regime.

Summary of the Invention

10

15

20

25

The present invention relates to use of compounds that antagonise the pharmacological actions of the endogenous neurokinins, particularly at the NK2 receptor, and compositions containing them, in treating the aforementioned disease conditions and, in particular, gastric asthma.

Brief Description of the Drawings

The accompanying drawings incorporated in and forming part of the specification illustrates principles of the invention. In the drawings:

Figure 1 is a graph representing the effect of tachykinin antagonism on esophageal-acid-induced bronchoconstriction; and

Figure 2 is a graph representing the effect of Example 1 on HCl infusion into allergic Guinea pigs.

Description

The present invention provides a method of treating gastric asthma in a patient in need thereof with an effective amount of an NK2 receptor antagonist.

In another aspect the present invention provides a method of treating adverse respiratory events associated with gastroesophageal reflux in a patient in need thereof with an effective amount of an NK2 receptor antagonist.

In yet another aspect the present invention provides the use of an NK2 receptor antagonist in a method of manufacture of a medicament for the treatment of gastric asthma.

In yet another aspect the present invention provides the use of an NK2 receptor antagonist in a method of manufacture of a medicament for the treatment of treating adverse respiratory events associated with gastroesophageal reflux.

In yet a further aspect the present invention provides an NK2 receptor antagonist for use in the treatment of gastric asthma or for the treatment of treating adverse respiratory events associated with gastroesophageal reflux.

Compounds with NK2 receptor antagonist properties are known in the scientific and patent literature, for example in WO9719060, EPA739891, WO9624582, EPA709376, EPA709375, EPA680962, WO9516682, WO9515961, WO9512577, WO9505377, EPA630887, EPA625509, WO9314084, WO9605193, WO9422822 and many other patent applications.

Preferred NK2 receptor antagonists are SR-48968, SR-144190, YM-38336, MEN-10627, MEN-11420, GR-159897, RPR-106145, PD-147714, FK-224, MDL-105212A, MDL-105172A, L-743986, S-16474 (all reviewed by von Sprecher et al (IDrugs, 73 (1998)) and the compounds of WO9512577.

A particularly preferred compound is (S)-N-[2-(3,4-dichlorophenyl)-4-[4-(2-oxoperhydropyrimidin-1-yl)piperidino]butyl]-N-methyl-benzamide of the formula (I) or a pharmaceutically acceptable salt thereof, such as the dihydrochloride.

25

5

10

20

which is disclosed in WO9512577.

-4-

NK2 receptor antagonists may be prepared according to the references or by standard literature methods. It is well known in the art how to test a compound to assess whether it is an NK2 receptor antagonist, for example see assays in the aforementioned patent applications such as in WO9512577.

In order to use an NK2 antagonist for the therapeutic treatment of gastric asthma (including prophylactic treatment) of mammals including humans, it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

5

10

15

20

25

30

Therefore, in another aspect, the present invention provides a method of treating gastric asthma in a patient in need thereof with a pharmaceutical composition comprising an effective amount of an NK2 receptor antagonist.

In a further aspect the present invention provides a pharmaceutical composition comprising an NK2 receptor antagonist for use in the treatment of gastric asthma.

Pharmaceutical compositions comprising an NK2 antagonist may be administered in standard manner, for example by oral, parenteral, buccal, nasal administration or by inhalation or insufflation. For these purposes the compounds of this invention may be formulated by means known in the art into the form of, for example, tablets, capsules, aqueous or oily solutions, suspensions, emulsions, gels, nasal sprays, finely divided powders or aerosols or nebulisers for inhalation, and for parenteral use (including intravenous, intramuscular or infusion) sterile aqueous or oily solutions or suspensions or sterile emulsions.

The pharmaceutical composition comprising an NK2 antagonist may also contain, or be co-administered (simultaneously or sequentially) with, one or more pharmacological agents of value in treating gastric asthma or a related or associated disease.

The pharmaceutical composition will normally be administered to humans so that, for example, a daily dose of 0.01 to 25 mg/kg body weight (and preferably of 0.1 to 5 mg/kg body weight) of compound is received. This daily dose may be given in divided doses as necessary, the precise amount of the compound received and the route of administration depending on the weight, age and sex of the patient being treated and on the particular disease condition being treated according to principles known in the art.

Typically, unit dosage forms will contain about 1 mg to 500 mg of a compound. For example a tablet or capsule for oral administration may conveniently contain up to 250 mg (and typically 5 to 100 mg) of a compound. In another example, for administration by inhalation, a compound may be administered in a daily dosage range of 5 to 100 mg, in a

-5-

single dose or divided into two to four daily doses. In a further example, for administration by intravenous or intramuscular injection or infusion, a sterile solution or suspension containing up to 10% w/w (and typically 5% w/w) of a compound.

5

10

15

20

25

30

It has been demonstrated that tracheal plasma extravasation, a feature of asthma, induced by esophageal acid can be prevented by blockade of NK1 receptors (Hamamoto et al., J. Appl. Physiol. 82/3: 738-745, 1997). In a particular aspect therefore, treatment of gastric asthma with an NK2 receptor antagonist may be augmented with an NK1 receptor antagonist; alternatively, gastric asthma may be treated with a single compound containing dual antagonist activity against both receptors. Therefore, in one aspect an NK2 antagonist may be co-administered with an NK1 antagonist with the advantage of blocking both neurogenic inflammation (plasma extravasation, edema, and mucous secretion) and bronchial hyperreactivity In another aspect the NK2 antagonist for use in this invention is also capable of antagonism at the NK1 receptors. That is, the compound for use in this invention is capable of antagonising the actions of both NK1 and NK2 receptors. Such compounds are known from the literature and may include compounds within the patent applications mentioned above as well as for example compounds within WO9610568, EPA733632, WO9637489, EPA699674, WO9722597, WO9628441, WO9827085, WO9834949, EPA776893, FRA2729952, FRA2729953, FRA2729954, WO9719926, WO9721680. WO9818761, WO9820010, WO28297, WO9634857, WO9634864, WO9639386, WO9827086, WO9714671, WO9898076694, WO9719942, WO9807722, WO9605193, WO9727188, WO9705110, EPA790248, EPA791592 and USP 5688960, 5696267, 5789422, 5691362, 5719147 and 5807865.

Therefore, in a further aspect, the present invention provides a method of treating gastric asthma in a patient in need thereof with an effective amount of an NK1/NK2 receptor antagonist.

In another aspect the present invention provides the use of an NK1/NK2 receptor antagonist in a method of manufacture of a medicament for the treatment of gastric asthma.

In yet a further aspect the present invention provides an NK1/NK2 receptor antagonist for use in the treatment of gastric asthma.

Gastroesophageal reflux, also known as heartburn, is a widespread disorder which affects millions of adults and children. Gastroesophageal reflux has been described as the retrograde movement of stomach contents into the esophagus. This is clearly discomforting.

-6-

If gastroesophageal reflux is not treated more severe disease conditions may occur such as esophagitis, ulceration, hemorrhage and related conditions. It has been estimated that approximately 40% of asthmatics suffer from gastroesophageal reflux. The aspiration of gastric contents into the airway or a vagal reflex arising from the esophagus may result in bronchoconstriction, as described by Harding and Richter, Chest. 111, 1359 (1997).

Gastroesophageal reflux is presently treated by antacids, histamine H₂-antagonists or proton pump inhibitors such as omeprazole. Relapse is common, regardless of therapy.

In another aspect the present invention provides an NK2 receptor antagonist for treating gastroesophageal reflux and in particular for the treatment of the adverse respiratory events such as chronic cough and asthma associated with gastroesophageal reflux.

The present invention is illustrated by the tests and results hereinbelow.

NK₁ and NK₂ In Vivo Functional Assay

10

15

20

25

30

The activity of a compound as an antagonist of NK1 and/or NK2 receptors may be demonstrated in vivo in laboratory animals as described in: Buckner et al. "Differential Blockade by Tachykinin NK1 and NK2 Receptor Antagonists of Bronchoconstriction Induced by Direct-Acting Agonists and the Indirect-Acting Mimetics Capsaicin, Serotonin and 2-Methyl-Serotonin in the Anesthetized Guinea Pig." J. Pharm. Exp. Ther., 1993, Vol 267(3), pp. 1168-1175. For experiments measuring the effects of infused acid into the esophagus on bronchoconstriction:

Compounds are tested in anesthetized guinea pigs pretreated with i.v. indomethacin (10 mg/kg, 20 min) and propranolol (0.5 mg/kg, 15 min). In some experiments thiorphan (10 mg/kg, 10 min), atropine (1 mg/kg, 25 min), 1-(\alpha-(2-isopropoxyphenyl)aceto)-3-(4-phenylquinuclidin-1-yl)-3R-(3,4-dichlorophenyl)piperidine (1 \mumol/kg, 25 min), Example I (various doses, 30 min) or their combination are also administered i.v. Saline (pH 5.5) or HCl (pH 0-1) is infused (0.5 ml over 1 min) into the esophagus. The infusate is confined to the esophagus due to ligations at the proximal and distal ends. Pulmonary resistance (a measure of bronchoconstriction) is monitored for at least thirty minutes after cessation of the infusion. Peak resistance values during the monitored period are then determined and expressed as a percentage of the baseline pulmonary resistance before the esophageal infusion.

For experiments measuring the effects of infused acid into the esophagus on tracheal plasma extravasation (edema):

-7-

Compounds are tested in anesthetized guinea pigs pretreated with i.v. 1-(α-(2-isopropoxyphenyl)aceto)-3-(4-phenylquinuclidin-1-yl)-3*R*-(3,4-dichlorophenyl)piperidine (3 μmol/kg, 25 min) or its vehicle, atropine (1 mg/kg, 25 min) indomethacin (10 mg/kg, 20 min), propranolol (0.5 mg/kg, 15 min), HOE-140 (0.1 μmol/kg, 12 min), thiorphan (10 mg/kg, 10 min) and Evans blue dye (30 mg/kg, 0 min). Saline (pH 5.5) or HCl (pH 0-1) is infused (0.5 ml over 1 min) into the esophagus. The infusate is confined to the esophagus due to ligations at the proximal and distal ends. Ten minutes following the esophageal infusion guinea pigs are perfused via the aorta (120 ml saline over 2 min) to remove intravascular dye. A section of trachea (1 cm) immediately caudal to the carina is removed, trimmed, weighed, and incubated in formamide for 18 hr at 60 °C. The dye content of the tissue is then determined by measuring Absorbance, at 620 nm, of the resulting formamide/Evans blue solution and converting this value into ng Evans blue/mg trachea via a standard curve.

Results of testing of (S)-N-[2-(3,4-dichlorophenyl)-4-[4-(2-oxoperhydropyrimidin-1-yl)piperidino]butyl]-N-methyl-benzamide (formula (I)) are as follows. 1-(α -(2-Isopropoxyphenyl)aceto)-3-(4-phenylquinuclidin-1-yl)-3R-(3,4-dichlorophenyl)piperidine is a known selective NK1 antagonist.

The graphs in Figures 1 and 2 clearly demonstrate that the NK2 receptor antagonist (Example I), either alone or in combination with the other treatments, is capable of completely inhibiting the increase in pulmonary resistance (i.e. bronchoconstriction) resulting from the infusion of 1 N HCl into the esophagus of guinea pigs (top graph). The muscarinic antagonist, atropine, or the NK1RA, 1-(α-(2-Isopropoxyphenyl)aceto)-3-(4-phenylquinuclidin-1-yl)-3*R*-(3,4-dichlorophenyl)piperidine, either alone or concomitantly have no effect on this increase in resistance (top graph). Even in an allergic, hyperreactive state (OA sensitized), as is often the case in asthma, NK2 receptor antagonism is able to anatagonize the adverse pulmonary effects of infused acid (bottom graph). This experimental data is representative of the effects of acid reflux in the esophagus of human asthmatics suffering from "gastric asthma" and illustrates the anticipated benefit of NK2 receptor antagonists in treating this syndrome.

Examples

15

20

25

30

The following illustrates representative pharmaceutical dosage forms which may be used for the therapeutic or prophylactic administration of NK1 antagonists or pharmaceutically-acceptable salt or in vivo hydrolysable ester thereof hereinafter referred to as 'Compound X'):

WO 00/25766

PCT/GB99/03536

-8-

	(:)	Table 1	
	(i)	Tablet 1	mg/tablet
		'Compound X'	100.0
		Lactose	77.5
		Providone	15.0
5		Croscarmellose sodium	12.0
		Microcrystalline cellulose	92.5
		Magnesium stearate	3.0
			300.0
10	(ii)	Tablet 2	mg/tablet
		'Compound X'	20.0
		Microcrystalline cellulose	410.0
		Starch	50.0
		Sodium starch glycolate	15.0
15		Magnesium stearate	5.0
			500.0
	(iii)	Capsule	mg/capsule
		'Compound X'	5.9
20		Lactose	392.9
		Sodium lauryl sulphate	1.2
			400.0
		•	
	(iv)	Capsule 2	mg/capsule
25		'Compound X'	29.6
		Lactose	331.4
		Sodium Lauryl Sulphate	1.0
			362.0
30	(v)	Injection 1 (1mg/mL)	mg/mL
		'Compound X' (free acid form)	1.0

		- 9-
	Dibasic sodium phosphate	12.0
	Monobasic sodium phosphate	0.7
	Sodium chloride	4.5
	1.0 N Sodium hydroxide solution	q.s
5	(pH adjustment to 7.0-7.5)	
	Water for injection	q.s. ad 1 mL

	(vi)	Injection 2 (10 mg/mL)	mg/mL
		'Compound X' (free acid form)	10.0
10		Monobasic sodium phosphate	0.3
		Dibasic sodium phosphate	1.1
		Polyethylene glycol 400	200.0
		0.1 N Sodium hydroxide solution	q.s.
		(pH adjustment to 7.0-7.5)	
15		Water for injection	q.s. ad 1 mL

(vii) Aerosol

'Compound X'

HFA 227 or HFA 134A with 5% ethanol

20

It will be appreciated that the above pharmaceutical compositions may be varied according to well-known pharmaceutical techniques to accommodate differing amounts and types of active ingredient 'Compound X'. The aerosol (vii) may be used in conjunction with a standard, metered dose aerosol dispenser.

PCT/GB99/03536

CLAIMS

- The use of an NK2 antagonist for treating or prevention of a condition in a patient in need thereof, comprising the step of administering a therapeutically-effective amount of the
 NK2 receptor antagonist, wherein the condition is gastric asthma or adverse respiratory events associated with gastroesophageal reflux.
 - 2. A use according to Claim 1 additionally comprising the step of administering a NK1 antagonist.

10

- 3. A use according to Claim 1 wherein the NK2 antagonist is also an NK1 antagonist.
- 4. A method according to any of Claim 1 or 2 wherein the NK2 antagonist has the structure

15

5. Use of an NK2 receptor antagonist for the manufacture of a medicament for the treatment or prevention of a condition, wherein the condition is gastric asthma or adverse respiratory events associated with gastroesophageal reflux.

20

- 6. A use according to any one of Claims 1, 2, 3 or 5 wherein the condition is gastric asthma.
- 7. A use according to any one of Claims 1, 2, 3 or 5 wherein the condition is adverse respiratory events associated with gastroesophageal reflux.
 - 8. An oral pharmaceutical composition for the treatment of a condition, the composition comprising:

-11-

a therapeutically-effective amount of an NK2 antagonist; and a diluent or carrier;


wherein the condition is gastric asthma or adverse respiratory events associated with gastroesophageal reflux.

5

- 9. A composition according to Claim 8 wherein the condition is gastric asthma.
- 10. A composition according to Claim 8 wherein the condition is adverse respiratory events associated with gastroesophageal reflux.

Example I 10 umoles/kg

Fig.1. 14007 ☐ Pooled saline infusion 1200 ☐1 N HCI **Pulmonary Resistance** <u>n=8</u> 1000 (%paseline) 800 600 400 200n=25 0 **Atropine** + 1mg/kg



SUBSTITUTE SHEET (RULE 26)

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



51) International Patent Classification 7: A61K 31/505, 31/00, 31/513,	A3	11) International Publication Number: WO 00/25766
A61P 11/06, 11/08	l	43) International Publication Date: 11 May 2000 (11.05.00
(22) International Application Number: PCT/GB99 (22) International Filing Date: 26 October 1999 (26) (30) Priority Data: 60/106,354 30 October 1998 (30.10.98) (71) Applicant (for all designated States except US): Z. LIMITED [GB/GB]; 15 Stanhope Gate, London W (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): LENGEL, Davi [US/US]; 1800 Concord Pike, P.O. Box 15437, Wilr DE 19850-5437 (US). RUMSEY, William, Leroy [1800 Concord Pike, P.O. Box 15437, Wilmingt 19850-5437 (US). (74) Agent: BRYANT, Tracey; Global Intellectual Prope traZeneca PLC, Mereside, Alderley Park, Macc Cheshire SK10 4TG (GB).	ENECTY 6L	KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RL SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AN AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. (88) Date of publication of the international search report: 21 September 2000 (21.09.00)
(54) Title: USE OF NK ANTAGONIST FOR TREATING	GAS	RIC ASTHMA
(57) Abstract		

with gastroesophageal reflux, including pharmaceu compositions.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
ВВ	Barbados	GH	Ghana	MG	Madagascar	ТJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
ВJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
) BR	Brazil	II.	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
l cc	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CI	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
l cu	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No PCT/GB 99/03536

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K31/505 A61K Ã61K31/00 A61P11/06 A61P11/08 A61K31/513 According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, CHEM ABS Data, BIOSIS, EMBASE. MEDLINE, SCISEARCH C. DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. Category WO 95 05377 A (ZENECA LTD) 7-10 X 23 February 1995 (1995-02-23) 1-6 Y example 18 claim 7 claims 1-9 page 1, line 26 - line 32 7-10 EP 0 791 592 A (PFIZER LTD ; PFIZER RES & X DEV (IE)) 27 August 1997 (1997-08-27) cited in the application 1-3,5,6page 2, line 8 - line 34 γ examples 1-9 -/--Further documents are listed in the continuation of box C. Patent family members are listed in annex. Х Special categories of cited documents: "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance invention "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other, such docucitation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or ments, such combination being obvious to a person skilled other means in the art. "P" document published prior to the international filling date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 07/07/2000 30 June 2000 Authorized officer Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040. Tx. 31 651 epo nl. Fax: (+31-70) 340-3016 Bonzano, C

5

INTERNATIONAL SEARCH REPORT

Inter. Idonal Application No PCT/GB 99/03536

.(Continu	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	····
ategory ·	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 12577 A (ZENECA LTD) 11 May 1995 (1995-05-11) cited in the application	7-10
'	page 1, paragraph 1 - paragraph 3 page 42; figures 1,1C claim 1	1-3,5,6
X	EP 0 625 509 A (ZENECA LTD) 23 November 1994 (1994-11-23)	7-10
Y	page 2, line 1 - line 30 	1-3,5,6
X	EP 0 630 887 A (ZENECA LTD) 28 December 1994 (1994-12-28)	7-10
Y	claims 1,9,11 	1-3,5,6
X	WO 97 27185 A (MARCHINGTON ALLAN PATRICK :MEADOWS SANDRA DORA (GB); MIDDLETON DON) 31 July 1997 (1997-07-31)	7-10
Υ	claim 1	1-3,5,6
Υ	US 5 697 112 A (COLAVITO BARBARA J ET AL) 16 December 1997 (1997-12-16) column 1, line 12	1-6
Y	MAYS: "Intrinsic asthma in adults" E.E.JAMA, vol. 236, no. 23, 1976, page 2626 XP000914287 page 2826, column 1 table 2	1-6
Y	KAVURU: "medical treatment" GASTROESOPHAGEAL REFLUX DISEASE AND AIRWAY DISEASE, - 1999 pages 179-203, XP000920806 page 181, paragraph 2	1-6

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims 1-3,5-10 relate to compounds defined by reference to a desirable pharmacological property, namely the activity as antagonist of NK1 or NK2 receptors.

The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is not fully possible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compounds by reference to their pharmacological profiles. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compounds structurally identified in claim 4, to those specifically disclosed in the description at page 3, lines 17-19, and to those described in the patents cited at page 3, lines 12-15 (as far as indexed by chemical abstracts), with due regard to the general idea underlying the present invention. Claims searched completely: 4. Claims searched incompletely: 1-3,5-10.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

information on patent family members .

Inter onal Application No PCT/GB 99/03536

Patent document cited in search repo	rt	Publication date		atent family nember(s)	Publication date
WO 9505377	Α	23-02-1995	AT AU CA CCZ DE EP ES HU JP NO NZ PK US US ZA	172971 T 677091 B 7388094 A 2166800 A 1129449 A 9600434 A 69414398 D 69414398 T 0714392 A 2123810 T 960727 A 75113 A 110672 A 9501439 T 960630 A 269575 A 313075 A 21096 A 5990130 A 5567700 A 5861392 A 9406142 A	15-11-1998 10-04-1997 14-03-1995 23-02-1995 21-08-1996 15-05-1996 10-12-1998 06-05-1999 05-06-1996 16-01-1999 16-02-1996 28-04-1997 20-06-1999 10-02-1997 16-02-1996 24-11-1997 27-05-1996 05-02-1997 23-11-1999 22-10-1996 19-01-1999 17-02-1995
EP 0791592	Α	27-08-1997	CA JP JP US US	2195923 A 2918508 B 9208579 A 6022971 A 5741790 A	28-07-1997 12-07-1999 12-08-1997 08-02-2000 21-04-1998
WO 9512577	A	11-05-1995	AT AU CA DE DE EP ES JP US	187963 T 8000594 A 2172377 A 69422303 D 69422303 T 0726893 A 2139760 T 9505036 T 5534525 A 5677317 A	15-01-2000 23-05-1995 11-05-1995 27-01-2000 18-05-2000 21-08-1996 16-02-2000 20-05-1997 09-07-1996 14-10-1997
EP 0625509	Α	23-11-1994	CA DE DE JP US	2123636 A 69404549 D 69404549 T 6340624 A 5521199 A	18-11-1994 04-09-1997 18-12-1997 13-12-1994 28-05-1996
EP 0630887	A	28-12-1994	AT AU CA CN DE ES FI GR HU IL	182583 T 673063 B 6320394 A 2124048 A 1098094 A 69419667 D 69419667 T 2135543 T 942381 A 3030875 T 70445 A,B 109734 A	15-08-1999 24-10-1996 15-12-1994 25-11-1994 01-02-1995 02-09-1999 06-04-2000 01-11-1999 25-11-1994 30-11-1999 30-10-1995 24-09-1998

INTERNATIONAL SEARCH REPORT

information on patent family members

Inter onal Application No
PCT/GB 99/03536

Patent document cited in search report		Publication date	ı	Patent family member(s)	Publication date
EP 0630887	A		IL JP NO NZ US	120895 A 6340625 A 941906 A 260566 A 5654299 A	24-09-1998 13-12-1994 25-11-1994 26-07-1996 05-08-1997
WO 9727185	A	31-07-1997	AU CA EP HR JP	1442297 A 2240964 A 0888337 A 970040 A 11505271 T	20-08-1997 31-07-1997 07-01-1999 30-06-1998 18-05-1999
US 5697112	Α	16-12-1997	NON	 :	